

Acute Effects of a Single, Oral dose of d9-tetrahydrocannabinol (THC) and Cannabidiol (CBD) Administration in Healthy Volunteers

R. Martín-Santos^{1,2,3*}, J.A. Crippa^{1, 4}, A. Batalla², S. Bhattacharyya¹, Z. Atakan¹, S. Borgwardt^{1,5}, P.Allen¹, M.Seal^{1,6}, K.Langohr^{3,7}, M. Farré³, AW.Zuardi⁴ and P.K. McGuire¹

¹Department of Psychosis Studies, Institute of Psychiatry, King's College London, UK; ²Department of Psychiatry, Institut Clínic de Neurociències, Hospital Clínic, IDIBAPS, CIBERSAM, Barcelona and Department of Psychiatry and Clinical Psychobiology, University of Barcelona, Spain; ³Human Pharmacology and Neurosciences, Hospital del Mar Research Institute - IMIM, Barcelona, Spain; ⁴Department of Neuroscience and Behavior, Faculty of Medicine of Ribeirão Preto, University of São Paulo, Brazil and INCT Translational Medicine, Ribeirão Preto, Brazil; ⁵Department of Psychiatry, UPK, University of Basel, Switzerland; ⁶Melbourne Neuropsychiatry Centre, The University of Melbourne, Australia; ⁷Department of Statistics and Operations Research, Polytechnic University of Catalonia, Barcelona, Spain

Abstract: *Rationale:* Animal and humans studies suggest that the two main constituents of *cannabis sativa*, delta-9-tetrahydrocannabinol (THC) and cannabidiol (CBD) have quite different acute effects. However, to date the two compounds have largely been studied separately.

Objective: To evaluate and compare the acute pharmacological effects of both THC and CBD in the same human volunteers.

Methods: A randomised, double-blind, cross-over, placebo controlled trial was conducted in 16 healthy male subjects. Oral THC 10 mg or CBD 600 mg or placebo was administered in three consecutive sessions, at one-month interval. Physiological measures and symptom ratings were assessed before, and at 1, 2 and 3 hours post drug administration. The area under the curve (AUC) between baseline and 3 hours, and the maximum absolute change from baseline at 2 hours were analysed by one-way repeated measures analysis of variance, with drug condition (THC or CBD or placebo) as the factor.

Results: Relative to both placebo and CBD, administration of THC was associated with anxiety, dysphoria, positive psychotic symptoms, physical and mental sedation, subjective intoxication (AUC and effect at 2 hours: $p < 0.01$), an increase in heart rate ($p < 0.05$). There were no differences between CBD and placebo on any symptomatic, physiological variable.

Conclusions: In healthy volunteers, THC has marked acute behavioural and physiological effects, whereas CBD has proven to be safe and well tolerated.

Keywords: Cannabis, Δ-9-THC-tetrahydrocannabinol, cannabidiol, unique dose, pharmacological acute effects, humans, induced anxiety, induced psychosis, review.

INTRODUCTION

Cannabis sativa preparations (marijuana, hashish, and others) are the illicit drugs most widely used in young people [1]. The plant has around 400 different chemical constituents, but two of its major psychoactive compounds are delta-9-tetrahydrocannabinol (THC) [2] and cannabidiol (CBD) [3,4].

THC acts as a partial agonist at specific endogenous cannabinoid receptors, termed CB1 and CB2, both members of the G-protein coupled receptor class [5]. The CB1 receptors are mainly expressed in the central nervous system, with a high density in the anterior cingulate, prefrontal cortex, medial temporal lobe and other areas [6] and are thought to mediate the majority of the effects of THC in the central nervous system. However, depending on the brain region, and whether the local CB1 receptors are expressed on neurons that release GABA or glutamate, THC can have either inhibitory or excitatory effects [7].

The acute administration of THC is associated with relaxation and enjoyment, but can also lead to unpleasant effects such as anxiety, psychotic symptoms, depression, apathy, and impairment of memory [8]. It has also been associated with impairments in

learning, motor coordination, slowed reaction time, impaired concentration during complex tasks, deficits in some executive functions, and impairments in some aspects of verbal processing, such as verbal fluency [9,10]. THC administration can also produce an increase in heart rate and orthostatic hypotension. However, the acute effects of THC and their time of onset are subject to wide inter-individual variation and due to differences in route of administration, rate of absorption, metabolism and the subject's expectation of its effects [11].

In contrast, CBD has a low affinity for CB1 receptors [12] and its molecular mechanism of action remains poorly understood. It may facilitate endocannabinoid signaling by inhibiting the cellular uptake and enzymatic hydrolysis of endocannabinoids [12]. It can also bind to CB1 and to serotonergic (5HT1A) receptors, inhibit adenosine uptake, and can activate vanilloid (TRPV1) receptors at micromolar concentrations [12-16]. CBD is pharmacologically active and can have anticonvulsant, sedative, anxiolytic [3,4,17,18] and antipsychotic effects [4, 19-25]. Unlike THC, CBD does not have acute effects on motor or cognitive performance [26, 27], nor does it have significant effects on pulse rate or blood pressure [28, 29]. Functional neuroimaging studies have confirmed the neurophysiological effects of THC and CBD are distinct and opposite [30-34]. Moreover, co-administration of CBD and THC may alter the pharmacological effect of the THC, in that CBD potentiates some of THC's desirable effects but attenuates some of its negative effects [29, 34-36]. However, it is difficult to establish which

*Address correspondence to this author at the Department of Psychiatry, Institut Clínic de Neurociències, Hospital Clínic, IDIBAPS, CIBERSAM, and Department of Psychiatry and Clinical Psychobiology, University of Barcelona, Villarroel, 170. 08036-Barcelona, Spain; Tel: 34 932275400; Fax: 34 932275538; E-mail: rmsantos@clinic.ub.es

CBD/THC ratios cause antagonism or potentiation, since other factors could interfere in the effects of these cannabinoids, such as the time between administrations of the two cannabinoids [37, 38]. Recent data showed the absence of significant differences between similar dose of oral THC and Sativex™, a plant extract with a 1: 1 proportion of both compound, on respect to subjective and physiological effects or pharmacokinetic [39, 40].

A better knowledge of the acute pharmacology effects of the two main compounds of the *cannabis sativa* may have implications for future research and therapeutics. We conducted a systematic review to assess the evidence for symptomatic and physiological effects of a single oral dose of THC and CBD in healthy volunteers. We reviewed literature in MEDLINE-PubMed database reporting studies with a cross-over, double-blind, placebo-controlled and randomised design in the last decade (2000-2011) (Table 1). We found nine studies which met our inclusion criteria in which seven studies compared THC to placebo [41-47], one with Modafinil [46], another with an active placebo (Diazepam) [48], and one in front morphine using an active placebo (Diazepam) [48]. Three of the studies had used cannabis extracts (with small proportion of CBD) [41, 44, 48] and one had compared CBD with placebo [49]. None of the studies had compared both compounds within the same sample.

Therefore we aimed to carry out a study with the objective of evaluating the acute effects of THC and CBD in the same group of healthy volunteers. Subjects were studied after a single dose of THC, CBD or placebo in three consecutive sessions separated by an interval of one month. Given the findings from previous studies [29, 50], our main hypothesis was that THC and CBD would have distinct effects on symptoms and physiological measures.

MATERIAL AND METHODS

Subjects

The study was conducted in accordance with the Declaration of Helsinki, approved by the local research committee (The Joint South London and Maudsley Trust and Institute of Psychiatry NHS Research Ethics Committee). All participants signed an informed consent form after full explanation of the study was given and were paid for their participation. Thirty right-handed, English-speaking healthy male volunteers, aged 18 to 42 years, were recruited through advertisement in local newspapers, posters and word-of-mouth referrals. Alcohol and illicit drug use was assessed in detail using a semi-structured questionnaire [51], and used to screen potential participants. Only individuals who had used cannabis less than 15 times in their lifetime and had not experienced any undesirable effects after use, such as anxiety and/or psychotic symptoms were included. They were also required not to have used cannabis in the previous month and abstain from using cannabis over the study duration. Exclusion criteria included those who had used any other psychotropic drug on a regular basis or drank more than 21 units of alcohol per week or had any psychiatric, neurological or severe medical illness history. Those with a family history of a psychotic illness were also excluded.

Sixteen right-handed male volunteers, with a mean (SD) age of 26.4 (5.3) years (range 20-42) were selected for the study. They had completed a mean (SD) of 16.46 (3.9) years of education. Nine subjects (56.3%) reported having used cannabis less than 5 times in their lifetime, while 7 (43.8%) reported having used cannabis on between 5-14 occasions. None had a history of substance abuse or dependence defined according to DSM-IV criteria, except for nicotine dependence. Seven subjects were current smokers, but only two subjects smoked more than 10 cigarettes/day. All subjects had Reading scores on the WRAT-R test [52] within the normal range (mean (SD) = 98.67 (7.078); range 79-108).

Participants remained under close clinical observation in the research centre for at least 3 hours after each administration, with this period extended if they had not yet completely recovered. All

participants agreed not to drive or use any machinery until the following day. A taxi was provided to take them home after each session.

Drugs

THC and CBD (approximately 99.6% and 99.9% pure, respectively) were supplied by THC-Pharm (Frankfurt, Germany) and STI Pharmaceuticals Ltd, (Brentwood, UK), and prepared by the Pharmacy Department of the Maudsley Hospital as identically appearing opaque capsules. The three drug conditions in the study were as follows: 10 mg THC, 600 mg CBD and placebo (flour). The doses of THC and CBD were selected on the basis of previous research [37,54-56] to produce a neurocognitive effect without provoking severe toxic, psychiatric or physical symptoms, which might confound interpretation of physiological and neuro-psychological data, or lead to the subject being unable to co-operate with the assessment.

Study Design

A crossover, double-blind, repeated measures design was used to compare the effects of THC, CBD and placebo. Participants were tested on three occasions at one-month intervals. The order of drug administration was pseudo-randomised to control for order effects. During the initial screening process, potential participants were familiarized with the testing procedures and questionnaires.

On each study day, subjects arrived at the research centre 1 hour before starting, having slept at least 6 hours and having had a standardised light breakfast. At each session, and before starting each assessment, urine samples were collected for screening for opiates, cocaine, amphetamines, benzodiazepines and THC using immunometric assay kits. None of the participants tested positive on any of the sessions. An indwelling intravenous catheter was then inserted into a subcutaneous vein in the forearm of the non-dominant arm. Thereafter, subjects remained seated in a quiet room throughout the session. Each drug was administered approximately after one hour of basal assessment.

Symptomatic Effects

Symptoms were evaluated at baseline and at 1, 2 and 3 hours after drug administration, using the Positive and Negative Psychotic Syndrome Scale (PANSS) [57], assessed by an experienced psychiatrist, and using a set of self-administered scales (below). The PANSS [57] a 30-item rating instrument was used to assess psychotic symptoms, with ratings based on a semi-structured clinical interview. Scores for each item range from 0 (absent) to 7 (extreme), and yield sub-scores for positive, negative, and general psychopathology domains. The self-administered scales comprised a 16-item version of the Visual Analogue Mood Scale (VAMS) [58], with four subscales: mental sedation or intellectual impairment, physical sedation or bodily impairments, anxiety effects and other types of feelings or attitudes. We also used the Addiction Research Centre Inventory (ARCI 49 item short form), a standardised measure of drug effects developed by Martin *et al* (1971) [59], comprising 49 true/false statements describing the subjective effects of various classes of substances. It has five empirically derived scales, measuring drug-induced euphoria (morphine-benzedrine group: MBG), stimulant-like effects (amphetamine group: A), intellectual efficiency and energy (benzedrine group: BG) and sedation (phenobarbital-chlorpromazine, alcohol group: PCAG), and dysphoria and somatic effects (lysergic acid: LSD). The Spielberger State Anxiety Inventory (STAI-T/S) [60] was used to assess state anxiety at hourly intervals, with subjects completing 20 items on current feelings and 20 on feelings in general.

Physiological Measures

Non-invasive systolic blood pressure (SBP), diastolic blood pressure (DBP), and heart rate were recorded at 1 hour before administration, immediately before drug administration (time 0, base-

Table 1. Systematic review (MEDLINE-PubMED, 2000-2011) of cross-over, double-blind, placebo-controlled, randomized studies of subjective and physiological effects of a single oral dose, THC, CBD, administration in healthy volunteers*.

Author (year)	Inclusion (In) /Exclusion (Ex)** criteria	M/F	M (SD) range	Drugs adminis- tered	Dose mg	Meas- ures hours	Clinical tools	Symptomatic effects		Physiological effects	Plasma concentrations ng/mL (Mean (SD) NA
								Increased	Decreased		
Sugarman <i>et al.</i> (2011) [46]	In: Healthy occasional volunteers THC+urine Ex: Any abuse/depend. Current psychi- atric disorder Physical illness	11/1	33.7 (7.7)	THC (dronabinol) +Placebo Modafinil +Placebo THC+Modafinil Placebo	15 +400 15+400	Basal, ½, 1, 1½, 2½, 3, 3½, 4, 4½ & 5h	ARCI DEQ POMS BP HR	THC ARCI (sedation, dysphoria) DEQ (“feel high”, “feel sedated”, & “feel the drug strength”)	THC POMS (vigor, depression) THC + Modafinil ARCI (euphoria)	THC HR increase Systolic BP low THC + Modafinil HR>increase	(SD) NA
Roser <i>et al.</i> (2008, 2009; Nadulski <i>et al.</i> , 2005 a,b) [41, 67- 69]	In: Healthy occasional volunteers Ex: Any abuse/depend. Current/past psychiatric disorder Positive urine analysis Pregnancy	12/12	27.9 (2.9) 18-45	THC Cannabis extract Placebo	10 THC:10 CBD:5.4	Basal, ½, 1, 1½, 2, 4, 7, 9 & 24h	AIR FTA NA	THC AIR (subjective level of intoxica- tion) Cannabis ext AIR Both similar	-	-	THC peak at 2h slightly > in F Similarly results with Cannabis ext. (THC and CBD).
Menetrey <i>et al.</i> (2005) Favrat <i>et al.</i> (2005) [42,70]	In: Healthy occasional volunteers Ex: Any abuse/depend. Current/past psychiatric disorder Physical illness	8/0	22-30	THC (dronabinol) Milk decoction Placebo	20 THC:16.5 THC:45.7 THC:1% CBD:0.4%	Basal, 1, 1½, 4, 5½, 7, 10 & 24h	VAS BP HR Conjun- tival redden- ing	THC & decoction VAS (strong feeling of high intoxication) > after the highest dose Decoction of 45.7 mg > Nausea and vomiting Two subjects excluded for anxiety (decoction 16.5 mg) and psychotic symptoms (dronabi- nol)	-	THC & decoc- tion HR slight/ moderate increased & conjunctival reddening	The highest mean THC was after ingestion the highest milk decoction.
Crippa <i>et al.</i> (2004) [49]	In: Healthy occasional volunteers Ex: Any abuse/depend. Personal/family current/past psychiatric disorder Physical illness Positive urine analysis	10/0	29.8 (5.1) 25-42	CBD Placebo	400	-½ (basal), 0, 1, & 1½ h	VAMS NA	CBD VAMS (mental sedation)	CBD VAMS (subjective anxiety)	-	NA

(Table 1) Contd....

Author (year)	Inclusion (In) /Exclusion (Ex)** criteria	M/F	M (SD) range	Drugs adminis- tered	Dose mg	Meas- ures hours	Clinical tools	Symptomatic effects		Physiologi- cal effects	Plasma concentra- tions ng/mL (Mean (SD))
								Increased	Decreased		
McDon- ald <i>et al.</i> (2003) [43]	In: Healthy occasional volunteers Ex: Any abuse/depend. Current/past psychiatric disorder Physical illness Low level education BMI: out of 19– 26 kg/m ² Positive urine analysis Pregnancy	18/19	23 (4.5) 18-45	THC (dronabinol) Placebo	7 15	Basal, 1/3, 11/3 & 21/3h	DEQ ARCI POMS BP HR	THC ARCI (stimulant- effects, marijuana- like effects, dysphoria, euphoria, somatic effects & sedation) DEG dose- dependently (“feel drug,” “feel high”, & “want more”) POMS dose- dependently (anxi- ety, fatigue, anger, & confusion)	THC ARCI (intellectual efficiency and energy)	THC HR increase dose depen- dently BP was not affected	NA
Wachtel <i>et al.</i> (2002) [44]	In: Healthy occasional volunteers Ex: Any abuse/depend. Current/past psychiatric disorder Physical illness Low level education BMI: out of 19– 26 kg/m ² Pregnancy	7/5	23 (4) 18-31	THC Whole-plant marijuana, Placebo	8.4 16.9 8.4 16.9	Basal, ½, 1, 1½, 2, 2½, 3, 4 & 5h	VAS DEQ POMS BP HR RR BT	THC dose- dependent DEQ, ARCI (mari- juana subscale & sedation) > mari- juana group THC-High condi- tion ARCI (stimulant effect, dysphoria & euphoria) > mari- juana group Marijuana DEQ & ARCI (marijuana scores and sedation) dose- dependently Marijuana-High condition VAS (sedated, drowsy and tired)	-	Any relevant physiological effect	THC increases dose dependent 1h after 11-OH-THC after 1.5h THC-High condi- tion > levels than marijuana-High condition
Curran <i>et al.</i> (2002) [45]	In: Healthy occasional volunteers Ex: Any abuse/depend Current psychi- atric disorder Physical illness Any drug use Positive urine analysis	15/0	24.2 (2.1) 18-30	THC (dronabinol) Placebo	7.5 15	Basal, 1, 2, 4, 6, 8, 24 & 48h	VAMS VAS NA	THC VAMS (drowsiness, anxiety) VAS (dizziness, dry mouth, palpitation and stoned feeling) No residual effects were found at 24h and 48h	THC VAMS (memory, concentra- tion)	THC HR increase on the high dose	THC peak at 2h after both high and low dose. 11-OH-THC levels same pattern. Levels at 24 & 48h were below limit detection

Data Analysis

Statistical analyses of these measures were carried out using SPSS (v.15) by two of the researchers (RMS and KL) blind to the drug conditions. The various measures obtained from the experimental sessions (symptomatic, physiological, and drug level data) were transformed to permit analysis of the differences in each variable relative to baseline. For each variable, the area under the curve (AUC) between baseline and 3 hours was calculated using the trapezoidal rule. The maximum absolute change from baseline at 2 hours was also determined. The AUC and the effect at 2 hours were analysed using a one-way repeated measures analysis of variance with drug condition (THC or CBD or placebo) as factor. When ANOVA showed significant effects for drug condition, post-hoc multiple comparisons were performed, using the Tukey's test for repeated measures. Correlations between whole blood levels of the drugs and its metabolites and statistical significant symptomatic effects, and physiological measures were analysed using Spearman's correlation coefficient. Differences associated with P-values lower than 0.05 were considered to be statistically significant. When necessary, Bonferroni multiple testing correction test was used.

RESULTS

Symptomatic Effects

Table 2 shows that there were highly significant differences between the effects of the THC in comparison to CBD and placebo. THC produced changes on positive and negative psychotic symptoms, and general psychopathology (PANSS), anxiety (STAI-S), dysphoria (ARCI), sedation (VAMS, ARCI), and the level of subjective intoxication (ASI, ARCI), as indexed by both the AUC and by the effect at 2 hours ($p < 0.001$). There was also difference on the VAMS anxiety ratings, which was significant at 2 hours ($p < 0.03$) between THC and CBD, but not in the AUC analysis. Some volunteers, 5 (33%) showed severe effects and became markedly paranoid and anxious, but there was a wide inter-subject variability, with a wide range of scores on the PANSS positive scale. Pair-wise comparisons revealed significant differences between the effects of THC relative to both placebo, and to CBD (Table 2). In contrast, there were no significant differences between the effects of CBD and placebo on any variable. The transient psychotic symptoms observed had resolved spontaneously within two hours. No psychopathological symptoms were reported on follow-up at next day, 1 and 3 weeks later.

(Figs. 1, 2, 3 and 4) show the effects of the drugs on each measure (ASI, STAI-S, VAMS, ARCI, and PANNS) at 1, 2, and 3 hours post administration.

PHYSIOLOGICAL EFFECTS AND PLASMATIC CONCENTRATIONS OF THC AND CBD

Physiological Parameters

There were significant differences between drug effects on heart rate (Table 3; Fig. 5). Pair-wise comparisons showed that this reflected an increase in heart rate with THC relative to both placebo, and to CBD (placebo vs. THC: $p = 0.0491$; THC vs. CBD: $p = 0.0133$; placebo vs. CBD: $p = 0.8596$). There was also a trend ($p < 0.07$) towards difference in the drug effects on diastolic blood pressure at 2 hours (Table 3).

Blood Levels

Mean (SD) whole blood levels of THC at 1, 2 and 3 hours after administration were 0.5 (0.8) ng/mL and 0.67 (0.66) ng/mL, and 0.44 (0.40) ng/mL, respectively. Mean (SD) whole blood levels of CBD at the same time points were 0.36 (0.64) ng/mL, 1.62 (2.98) ng/mL and 3.4 (6.42) ng/mL, respectively. Levels of 11-OH-THC and THC-COOH were elevated after administration of THC (but not CBD or placebo) and followed a similar time course (Fig. 6).

Relationship between Blood Levels and Acute Symptomatic Effects

Both the level of subjective intoxication (ASI) and the PANSS total score (PANSS-TS) were directly correlated with THC-COOH levels at 1 hour post drug administration ($\rho = 0.665$; $p = 0.009$; $\rho = 0.687$; $p = 0.007$), and with THC levels at 3 hours post drug administration ($\rho = 0.760$; $p = 0.002$; $\rho = 0.731$; $p = 0.003$). Negative symptom levels (PANSS-N) also showed a positive correlation with both THC and 11-OH-THC levels at 3 hours post drug administration ($\rho = 0.813$; $p < 0.001$; $\rho = 0.727$; $p = 0.003$). We did not find significant correlation between heart rate and neither THC, 11-OH-THC nor THC-COOH whole blood levels.

DISCUSSION

Acute Symptomatic Effects

The administration of a single oral dose of THC produced the typical transient effects previously described for this substance in an experimental laboratory setting: feelings of anxiety, euphoria, dysphoria and subjective intoxication. Positive and negative psychotic symptoms were also evident in some, but not all subjects, again consistent with previous studies [63-65]. In the review done, seven of nine studies described "feel high", dysphoria, and subjective intoxication [41,42-44,46,48,66-69] (Table 1). The intensity of symptomatology appeared to be dose-dependent [42, 44]. Moreover, from the 146 subjects involved in the review, 3 (2.1%) were excluded because they presented severe acute psychotic symptomatology during the study [42, 58, 70]. In our study, 5 (33%) subjects presented transient psychotic symptomatology in the THC session, which resolved spontaneously in two hours. This variability probably reflects differences in individual, or genetic susceptibility to THC proneness to psychosis [71, 72].

Although studies in both experimental animals [73-77] and healthy volunteers [18, 29, 34, 49,78,79] have shown that CBD has anxiolytic properties, there were remarkably few differences between the effects of CBD and placebo on anxiety [17], save for a reduction in the VAMS anxiety scale at 2 hours post administration. However, in such previous human studies, the anxiolytic effect of CBD has only been evident in subjects in whom anxiety had already been induced experimentally, in contrast to the subjects in the present study. In addition, in animal models, the effect of CBD on anxiety appears to follow an inverted U-shaped dose-response curve [4, 75]. The dose of CBD used in the present study was higher than in previous human anxiety experimental studies (60-300 mg/day), [18, 29, 34, 49,78] and so may have exceeded the dose associated with a clear anxiolytic effect. Unlike THC, CBD had no effects on sedation, intoxication, mood or psychotic symptoms. These data suggest that CBD alone has remarkably few symptomatic effects in non-anxious healthy subjects, which is important in relation to the potential therapeutic utility of CBD in neurology, psychiatry and other fields of medicine [4, 24]. Recently, a double-blind, randomised study showed that CBD reduces anxiety induced by a simulation public speaking test in a group of patients with generalized social anxiety disorder to a similar response as healthy controls [79].

Physiological Measures

THC increased the heart rate as observed in other studies [42, 43, 45-48], but did not produce an increased systolic and diastolic blood pressure or an orthostatic hypotension, although there was a tendency for an effect on diastolic blood pressure [11]. This may reflect an effect of the THC mediated by sympathetic activation and cholinergic inhibition [80]. As expected from previous investigations [28, 29], CBD did not have any significant physiological effects.

Table 2. Results of Symptomatic Effects Comparisons after a Single Oral dose of Placebo, THC, CBD and Placebo Administration with Respect to the Area Under the Curve (AUC) and Effect at 2 Hours

	AUC				Effect at 2 hours			
	F	p		p*	F	p		p*
Symptomatic effects								
ASI	7.81	0.002	1	<0.001	14.33	<0.001	1	<0.001
			2	0.929			2	0.778
			3	0.003			3	<0.001
STAI-S	6.20	0.006	1	0.002	10.50	0.001	1	<0.001
			2	0.455			2	0.354
			3	0.055			3	0.005
VAMS								
Anxiety	2.46	0.105			3.97	0.03	1	0.179
							2	0.634
							3	0.020
Mental sedation	4.67	0.018	1	0.010	6.89	0.004	1	0.001
			2	0.517			2	0.739
			3	0.166			3	0.015
Physical sedation	3.67	0.039	1	0.019	6.18	0.006	1	0.002
			2	0.374			2	0.417
			3	0.358			3	0.084
Other feelings	0.45	0.64			0.20	0.816		
ARCI								
Stimulant-like effects-A	2.42	0.111			2.86	0.076		
Euphoria-MBG	2.22	0.314			2.73	0.084		
Dysphoria-LSD	9.16	0.001	1	0.001	15.03	0.001	1	<0.001
			2	0.963			2	0.535
			3	<0.001			3	<0.001
Intellectual efficiency-BG	4.76	0.019	1	0.024	2.85	0.077		
			2	0.996				
			3	0.023				
Sedation-PCAG	8.33	0.002	1	<0.001	11.32	<0.001	1	<0.001
			2	0.928			2	0.845
			3	0.003			3	<0.001
PANNS								
General psychopathology	9.10	<0.001	1	<0.001	10.71	<0.001	1	<0.001
			2	0.668			2	0.91
			3	0.003			3	<0.001
Positive symptoms	9.14	0.001	1	<0.001	5.37	0.010	1	0.010
			2	0.966			2	0.975
			3	<0.001			3	0.019
Negative symptoms	5.65	0.008	1	0.002	5.73	<0.001	1	0.002
			2	0.359			2	0.317
			3	0.109			3	0.131

ASI= Subjective level of intoxication; STAI-S= Spielberger State Anxiety Inventory; VAMS= Visual Analogue Mood Scale; ARCI= Addiction Research Center Inventory; PANNS= Positive and Negative Psychotic Symptomatology Scale

*Pair wise comparisons: 1) placebo vs. THC, 2) placebo vs. CBD, and 3) THC vs. CBD




Fig. (1). Changes from baseline over time in the level of subjective intoxication (ASI) score and the level of anxiety (STAI-S) after oral administration of 10 mg THC, 600 mg CBD, and placebo. The figure shows mean ($\pm$ SEM) values.




Fig. (2). Changes from baseline over time in anxiety level, mental and physical sedation and other feelings (VAMS) after administration of 10 mg THC, 600 mg CBD, and placebo. The figure shows mean ($\pm$ SEM) values.




Fig. (3). Changes in subjective symptomatology related to drug intoxication: stimulant effects, induced euphoria, dysphoria, intellectual efficiency and sedation (ARCI) scores after administration of 10 mg THC, 600 mg CBD, and placebo. The figure shows mean ($\pm$ SEM) values




Fig. (4). Changes from baseline over time in positive and negative psychotic symptomatology and total score of general psychopathology of PANSS after administration of THC, CBD, and placebo. The figure shows mean ($\pm$ SEM) values.




Fig. (5). Changes from baseline over time in heart rate after oral administration of 10 mg of THC, 600 mg of CBD, and placebo. Figure shows mean ($\pm$ SEM) values.




Fig. (6). Time course of THC, 11-OH-THC and THC-COOH whole blood levels after oral administration of 10 mg of THC, 600 mg of CBD, and placebo. Figure shows mean ($\pm$ SEM) values.

Table 3. Results of Physiological Effects Comparisons after a Single Oral dose of Placebo, THC, CBD and Placebo Administration with Respect to the Area Under the Curve (AUC) and Effect at 2 Hours

Physiological parameters	AUC				Effect at 2 hours			
	F	p		p*	F	p		p*
Systolic blood pressure	0.96	0.397			1.17	0.327		
Diastolic blood pressure	0.27	0.769			2.44	0.07		
Heart rate	4.72	0.019	1	0.010	4.83	0.016	1	0.049
			2	0.924			2	0.859
			3	0.037			3	0.013

*Pair wise comparisons: 1) placebo vs. THC, 2) placebo vs. CBD, and 3) THC vs. CBD

Whole Blood Drug Concentration Levels

Although some previous studies have reported that THC plasma concentrations were out of phase with its behavioural, cognitive or endocrine effects [61,62, 81, 82], we found that the level of subjective intoxication (ASI) and the severity of positive and negative total score (PANSS-TS) correlated with whole blood levels of 11-OH-THC at 1 hour post drug administration, and with the levels of THC at 3 hours post drug administration.

Limitations

Some methodological limitations of this study need to be noted. First, we used a within-subject cross-over design, which minimised the confounding of effects of inter-subject differences, but was logistically demanding, limited the total number of participants that could be studied. In an effort to minimise the potentially confounding effects of previous substance use, we restricted inclusion to volunteers who has taken cannabis less than 15 times in their life-

time, with none in the last month. However, for ethical reasons, it was not possible to study participants who were completely cannabis naïve. The subjective effects of cannabis may be greater at the first time of use [11, 17], so we might have observed different results in a sample with more experience with cannabis. In the systematic review we observed that one of the three subjects, a woman, who presented acute psychotic symptoms was from a study in naïve subjects [48] (Table 1). The dose of THC chosen for this study (10mg) was designed to be comparable to that delivered from a typical cannabis cigarette, and it is possible that had we used a higher dose, effects on cognitive performance may have been more evident.

In summary, the data from the present study suggest that a single dose of THC, comparable to that delivered from a cannabis cigarette, had significant acute symptomatic and physiological effects in healthy volunteers. Moreover, CBD has confirmed to be safe and well-tolerated in humans as previously observed [25].

CONFLICTS OF INTEREST

The authors confirm that this article content has no conflicts of interest.

ACKNOWLEDGEMENTS

This study was supported by grants from the Psychiatry Research Trust, UK and the Ministerio de Sanidad y Consumo: Plan Nacional sobre Drogas: 2006I101, and PI041731/2011, and the Generalitat de Catalunya: SGR2009/1435, Spain. JAC and AWZ are recipients of a CNPq Productivity fellowship (Brazil). SB has received support from the Medical Research Council, UK (Joint MRC/Priority Clinical research training fellowship; G0501775) and is currently supported by a NIHR Clinician Scientist Award (NIHR CS-11-001).

REFERENCES

- [1] European Monitoring Centre for Drugs and Drug Addiction (EM-CDDA) Annual report 2011: the state of the drugs problem in Europe. Luxembourg: Publications Office of the European Union 2011.
- [2] Gaoni Y, Mechoulam R. The isolation of delta-1-tetrahydrocannabinol and other neutral cannabinoids from hashish. *J Am Chem Soc* 1971; 93: 217-24.
- [3] Mechoulam R, Hanus L. Cannabidiol: an overview of some chemical and pharmacological aspects. Part I: chemical aspects. *Chem Phys Lipids* 2002; 121: 35-43.
- [4] Crippa JAS, Zuardi AW, Hallak JEC. Therapeutic use of the cannabinoids in psychiatry. *Rev Bras Psiquiatr* 2012; 32 (Suppl 1): 556-65.
- [5] Grotenhermen F. Cannabinoids. *Curr drug targets CNS Neurol Disor* 2005; 4: 507-30.
- [6] Iversen L. Cannabis and the brain. *Brain* 2003; 126: 1252-70.
- [7] Pertwee RG. The diverse CB1 and CB2 receptor pharmacology of three plant cannabinoids: Delta9-tetrahydrocannabinol, cannabidiol and Delta-9-tetrahydrocannabivarin. *Br J Pharmacol* 2008; 153: 199-215.
- [8] Devane WA, Dysarz FA, Johnson MR, Melvin LS, Howlett AC. Determination and characterization of a cannabinoid receptor in rat brain. *Mol Pharmacol* 1988; 34: 605-13.
- [9] Solowij N. Cannabis and cognitive functioning. Cambridge: Cambridge University Press 1998.
- [10] Kelleher LM, Stough C, Sergejew AA, Rolfe T. The effects of cannabis on information-processing speed. *Addict Behav* 2004; 29: 1213-9.
- [11] Gorelick DA, Heishman SJ. Methods for clinical research involving cannabis administration. *Methods Mol Med* 2006; 123: 235-53.
- [12] Bisogno T, Hanus L, De Petrocellis L, et al. Molecular targets for cannabidiol and its synthetic analogues: effect on vanilloid VR1 receptors and on the cellular uptake and enzymatic hydrolysis of anandamide. *Br J Pharmacol* 2001; 134: 845-52.
- [13] Russo EB, Burnett A, Hall B, Parker KK. Agonistic properties of cannabidiol at 5-HT1a receptors. *Neurochem Res* 2005; 30: 1037-43.
- [14] Carrier EJ, Auchampach JA, Hillard CJ. Inhibition of an equilibrative nucleoside transporter by cannabidiol: a mechanism of cannabinoid immunosuppression. *Proc Natl Acad Sci U S A* 2006; 103: 7895-900.
- [15] Thomas R, Cheshier G. The pharmacology of marijuana. *Med J Aust* 1973; 2: 229-37.
- [16] Soares VDEP, Campos AC, Bortoli VC, Zangrossi H Jr, Huimaraes FS, Zuardi AW. Intra-dorsal periaqueductal gray administration of cannabidiol blocks panic-like response by activating 5-HT1A receptors. *Behav Brain Res* 2010; 213: 225-9.
- [17] Crippa JA, Zuardi AW, Martin-Santos R, et al. Cannabis and anxiety: a critical review of the evidence. *Hum Psychopharmacol* 2009; 24: 513-23.
- [18] Crippa JA, Darenusson GN, Ferrari TB, et al. Neural basis of anxiolytic effects of cannabidiol (CBD) in generalized social anxiety disorder: a preliminary report. *J Psychopharmacol* 2011; 25: 121-30.
- [19] Leweke FM, Schneider U, Radwan M, Schmidt E, Emrich HM. Different effects of nabilone and cannabidiol on binocular depth inversion in Man. *Pharmacol Biochem Behav* 2000; 66: 175-81.
- [20] Krystal JH, D'Souza DC, Madonick S, Petrakis IL. Toward a rational pharmacotherapy of comorbid substance abuse in schizophrenic patients. *Schizophr Res* 2000; 35 S35-49.
- [21] Zuardi AW, Morais SL, Guimaraes FS, Mechoulam R. Antipsychotic effect of cannabidiol. *J Clin Psychiatry* 1995; 56: 485-6.
- [22] Zuardi AW, Crippa JA, Hallak JE, Moreira FA, Guimarães FS. Cannabidiol, a Cannabis sativa constituent, as an antipsychotic drug. *Braz J Med Biol Res* 2006 (a); 39: 421-9.
- [23] Zuardi AW, Hallak JE, Dursun SM, et al. Cannabidiol monotherapy for treatment-resistant schizophrenia. *J Psychopharmacol* 2006(b); 20: 683-6.
- [24] Zuardi AW. Cannabidiol: from an inactive cannabinoid to a drug with wide spectrum of action. *Rev Bras Psiquiatr* 2008; 30: 271-80.
- [25] Bergamaschi MM, Queiroz RH, Zuardi AW, Crippa JA. Safety and Side Effects of Cannabidiol, a Cannabis sativa Constituent. *Curr Drug Saf* 2011; 6: 237-49.
- [26] Consroe P, Carlini EA, Zwicker AP, Lacerda LA. Interaction of cannabidiol and alcohol in humans. *Psychopharmacology (Berl)* 1979; 66: 45-50.
- [27] Belgrave BE, Bird KD, Cheshier GB, et al. The effect of cannabidiol, alone and in combination with ethanol, on human performance. *Psychopharmacology (Berl)* 1979; 64: 243-6.
- [28] Karniol IG, Carlini EA. Pharmacological interaction between cannabidiol and delta 9-tetrahydrocannabinol. *Psychopharmacologia* 1973; 33: 53-70.
- [29] Zuardi AW, Shirakawa I, Finkelfarb E, Karniol IG. Action of cannabidiol on the anxiety and other effects produced by delta 9-THC in normal subjects. *Psychopharmacology (Berl)* 1982; 76: 245.
- [30] Fusar-Poli P, Crippa JA, Bhattacharyya S, et al. Distinct effects of [delta]9-tetrahydrocannabinol and cannabidiol on neural activation during emotional processing. *Arch Gen Psychiatry* 2009; 66: 95-105.
- [31] Fusar-Poli P, Allen P, Bhattacharyya S, et al. Modulation of effective connectivity during emotional processing by Delta9-tetrahydrocannabinol and cannabidiol. *Int J Neuropsychopharmacol* 2010; 13: 421-32.
- [32] Bhattacharyya S, Morrison PD, Fusar-Poli P, et al. Opposite effects of delta-9-tetrahydrocannabinol and cannabidiol on human brain function and psychopathology. *Neuropsychopharmacology* 2010; 35: 764-74.
- [33] Borgwardt SJ, Allen P, Bhattacharyya S, et al. Neural basis of Delta-9-tetrahydrocannabinol and cannabidiol effects during response inhibition. *Biol Psychiatry* 2008; 64: 966-73.
- [34] Karniol IG, Shirakawa I, Kasinski N, Pfeifferman A, Carlini EA. Cannabidiol interferes with the effects of delta 9 - tetrahydrocannabinol in man. *Eur J Pharmacol* 1974; 28: 172-7.
- [35] Dalton WS, Martz R, Lemberger L, Rodda BE, Forney RB. Influence of cannabidiol on delta-9-tetrahydrocannabinol effects. *Clin Pharmacol Ther* 1976; 19: 300-9.
- [36] Russo E, Guy GW. A tale of two cannabinoids: the therapeutic rationales for combining tetrahydrocannabinol and cannabidiol. *Med Hypotheses* 2006; 66: 234-46.
- [37] Zuardi AW, Crippa JA, Hallak JE. Cannabis sativa: the plant that can induce unwanted effects and also treat them. *Rev Bras Psiquiatr* 2011; 32 Suppl 1: S1-2.

- [38] Zuardi AW, Hallak JEC, Crippa JAS. Interaction between cannabidiol (CBD) and delta-9-tetrahydrocannabinol (THC): influence of administration interval and dose ratio between the cannabinoids. *Psychopharmacology* 2012; 219: 247-9.
- [39] Karschner EL, Darwin WD, McMahon RP, *et al.* Subjective and physiological effects after controlled Sativex and oral THC administration. *Clin Pharmacol Ther* 2011; 89: 400-7.
- [40] Karschner EL, Darwin WD, Goodwin RS, Wright S, Huestis MA. Plasma cannabinoid pharmacokinetics following controlled oral delta-9-tetrahydrocannabinol and oromucosal cannabis extract administration. *Clin Chem* 2011; 57: 66-75.
- [41] Roser P, Juckel G, Rentzsch J, Nadulski T, Gallinat J, Stadelmann AM. Effects of acute oral Delta-9-tetrahydrocannabinol and standardised cannabis extract on the auditory P300 event-related potential in healthy volunteers. *Eur Neuropsychopharmacol* 2008; 18: 569-77.
- [42] Ménétrey A, Augsburger M, Favrat B, *et al.* Assessment of driving capability through the use of clinical and psychomotor tests in relation to blood cannabinoids levels following oral administration of 20 mg dronabinol or of a cannabis decoction made with 20 or 60 mg Delta-9-THC. *J Anal Toxicol* 2005; 29: 327-38.
- [43] McDonald J, Schleifer L, Richards JB, de Wit H. Effects of THC on behavioral measures of impulsivity in humans. *Neuropsychopharmacology* 2003; 28: 1356-65.
- [44] Wachtel SR, ElSohly MA, Ross SA, Ambre J, de Witt H. Comparison of the subjective effects of Delta(9)-tetrahydrocannabinol and marijuana in humans. *Psychopharmacology (Berl)* 2002; 161: 331-39.
- [45] Curran HV, Brignell C, Fletcher S, Middleton P, Henry J. Cognitive and subjective dose-response effects of acute oral Delta-9-tetrahydrocannabinol (THC) in infrequent cannabis users. *Psychopharmacology (Berl)* 2002; 164: 61-70.
- [46] Sugarman, DE, Poling J, Sofuoglu M. The safety of modafinil in combination with oral 9-tetrahydrocannabinol in humans. *Pharmacol Biochem Behav* 2011; 98: 94-100.
- [47] Kaufmann RM, Kraft B, Frey R, *et al.* Acute psychotropic effects of oral cannabis extract with a defined content of Delta-9-tetrahydrocannabinol (THC) in healthy volunteers. *Pharmacopsychiatry* 2010; 43: 24-32.
- [48] Naef M, Curatolo M, Petersen-Feliz S, Arendt-Nielsen L, Zbinden A, Brenneisen R. The analgesic effect of oral delta-9-tetrahydrocannabinol (THC), morphine, and a THC-morphine combination in healthy subjects under experimental pain conditions. *Pain* 2003; 105: 79-88.
- [49] Crippa JA, Zuardi AW, Garrido GE, *et al.* Effects of cannabidiol (CBD) on regional cerebral blood flow. *Neuropsychopharmacology* 2004; 29: 417-26.
- [50] Morgan CJ, Curran HV. Effects of cannabidiol on schizophrenia-like symptoms in people who use cannabis. *Br J Psychiatry* 2008; 192: 306-7.
- [51] McLellan AT, Kushner H, Metzger D, *et al.* The Fifth Edition of the Addiction Severity Index. *J Subst Abuse Treat* 1992; 9: 199-13.
- [52] Nelson HE. National Adult Reading Test (NART). Test Manual. NFER-Nelson: Windsor, Berkshire, England 1982.
- [53] Agurell S, Carlsson S, Lindgren JE, Ohlsson A, Gillespie H, Hollister L. Interactions of delta-9-tetrahydrocannabinol with cannabidiol and cannabidiol following oral administration in man. Assay of cannabidiol and cannabidiol by mass fragmentography. *Experientia* 1981; 37: 1090-2.
- [54] Cheshier GB, Bird KD, Jackson DM, Perrignon A, Starmer GA. The effects of orally administered delta-9-tetrahydrocannabinol in man on mood and performance measures: a dose response study. *Pharmacol Biochem Behav* 1990; 35: 861-4.
- [55] Lewke FM, Schneider U, Thies M, Munte TF, Emrich HM. Effects of synthetic delta-9 tetrahydrocannabinol on binocular depth inversion of natural and artificial objects in man. *Psychopharmacology (Berl)* 1999; 142: 230-5.
- [56] Koethe D, Gerth CW, Neatby MA, *et al.* Disturbances of visual information processing in early states of psychosis and experimental delta-9-tetrahydrocannabinol altered states of consciousness. *Schizophr Res* 2006; 88: 142-50.
- [57] Kay SR, Fiszbein A, Opler LA. The Positive and Negative Syndrome Scale (PANSS) for schizophrenia. *Schizophr Bull* 1987; 13: 261-76.
- [58] Folstein MF, Luria R. Reliability, validity, and clinical application of the Visual Analogue Mood Scale. *Psychol Med* 1973; 3: 479-86.
- [59] Martin WR, Sloan JW, Sapira JD, Jasinski DR. Physiologic, subjective, and behavioral effects of amphetamine, methamphetamine, ephedrine, phenmetrazine, and methylphenidate in man. *Clin Pharmacol Ther* 1971; 12: 245-58.
- [60] Spielberger C. Manual of the State-Trait Anxiety Inventory. CA Consulting, Palo Alto: Psychologists Press Inc. 1983.
- [61] Huestis MA, Henningfield JE, Cone EJ. Blood cannabinoids. II. Models for the prediction of time of marijuana exposure from plasma concentrations of delta-9-tetrahydrocannabinol (THC) and 11-nor-9-carboxy-delta-9-tetrahydrocannabinol (THCCOOH). *J Anal Toxicol* 1992; 16: 283-90.
- [62] Huestis MA. Human cannabinoid pharmacokinetics. *Chem Biodivers* 2007; 4: 1770-804.
- [63] D'Souza DC, Perry E, MacDougall L, *et al.* The psychotomimetic effects of intravenous delta-9-tetrahydrocannabinol in healthy individuals: implications for psychosis. *Neuropsychopharmacology* 2004; 29: 1558-72.
- [64] Morrison PD, Zois V, McKeown DA, *et al.* The acute effects of synthetic intravenous D9-tetrahydrocannabinol on psychosis, mood and cognitive functioning. *Psychol Med* 2009; 1: 1-10.
- [65] Sewell RA, Skosnik PD, Garcia-Sosa I, Ranganathan M, D'Souza DC. Behavioral, cognitive and psychophysiological effects of cannabinoids: relevance to psychosis and schizophrenia. *Rev Bras Psiquiatr* 2011; 32 (Suppl 1): S15-30.
- [66] Kraft B, Frickey NA, Kaufmann RM, *et al.* Lack of analgesia by oral standardised cannabis extract on acute inflammatory pain and hyperalgesia in volunteers. *Anesthesiology* 2008; 109: 101-10.
- [67] Roser P, Gallinat J, Weinberg G, Juckel G, Gorynia I, Stadelmann AM. Psychomotor performance in relation to acute oral administration of Delta-9-tetrahydrocannabinol and standardised cannabis extract in healthy human subjects. *Eur Arch Psychiatry Clin Neurosci* 2009; 259: 284-92.
- [68] Nadulski T, Sporkert F, Schnelle M, *et al.* Simultaneous and sensitive analysis of THC, 11-OH-THC, THC-COOH, CBD, and CBD by GS-MS in plasma after oral application of small doses of THC and cannabis extract. *J Anal Toxicol* 2005; 29: 782-9.
- [69] Nadulski T, Pragst F, Weinberg G, *et al.* Randomised, double-blind, placebo-controlled study about the effects of cannabidiol (CBD) on the pharmacokinetics of delta-9-tetrahydrocannabinol (THC) after oral application of THC versus standardised cannabis extract. *Ther Drug Monit* 2005; 27: 799-810.
- [70] Favrat B, Menetrey A, Augsburger M, *et al.* Two cases of cannabis acute psychosis following the administration of oral cannabis. *BMC Psychiatry* 2005; 5: 17.
- [71] Luzzi S, Morrison PD, Powell J, di Forti M, Murray RM. What is the mechanism whereby cannabis use increases risk of psychosis? *Neurotox Res* 2008; 14: 105-12.
- [72] Bhattacharyya S, Atakan Z, Martín-Santos R, *et al.* Preliminary report of biological basis of sensitivity to the effects of cannabis on psychosis: AKT1 and DAT1 genotype modulates the effects of delta-9-tetrahydrocannabinol on midbrain and striatal function. *Mol Psychiatry* 2012; doi: 10.1038/mp.2011.187.
- [73] Zuardi AW, Karniol IG. Changes in the conditioned emotional response of rats induced by delta-9-THC, CBD and mixture of the two cannabinoids. *Arq Biol Tecnol* 1983; 26: 391-7.
- [74] Musty RE, Conti LH, Mechoulam R. Anxiolytic properties of cannabidiol. In: Harvey DJ Ed. *Marihuana '84. Proceedings of the Oxford Symposium on Cannabis*. Oxford, UK: IRL Press Limited, 1984: pp. 713-9.
- [75] Guimarães FS, Chiaretti TM, Graeff FG, Zuardi AW. Antianxiety effect of cannabidiol in the elevated plus-maze. *Psychopharmacology* 1990; 100: 558-9.
- [76] Onaivi ES, Green MR, Martin BR. Pharmacological characterization of cannabinoids in the elevated plus maze. *J Pharmacol Exp Ther* 1990; 253: 1002-9.
- [77] Moreira FA, Aguiar DC, Guimarães FS. Anxiolytic-like effect of cannabidiol in the rat Vogel conflict test. *Prog Neuropsychopharmacol Biol Psychiatry* 2006; 30: 1466-71.
- [78] Zuardi AW, Cosme RA, Graeff FG, Guimarães FS. Effects of ipsapirone and cannabidiol on human experimental anxiety. *J Psychopharmacol* 1993; 7: 82-8.
- [79] Bergamaschi MM, Costa Queiroz RH, Hortes M, *et al.* Cannabidiol reduces the anxiety induced by simulated public speaking in treatment-naïve social phobia patients. *Neuropsychopharmacology* 2011; 36: 1219-26.

- [80] Jones RT. Cardiovascular system effects of marijuana. *J Clin Pharmacol* 2002; 42: 585-635.
- [81] Cocchetto DM, Owens SM, Perez-Reyes M, DiGiuseppi S, Miller LL. Relationship between plasma delta-9-tetrahydrocannabinol concentration and pharmacologic effects in man. *Psychopharmacology (Berl)* 1981; 75: 158-64.
- [82] Cone EJ, Huestis MA. Relating blood concentrations of tetrahydrocannabinol and metabolites to pharmacologic effects and time of marijuana usage. *Ther Drug Monit* 1993; 15: 527-32.

Received: March 30, 2012

Accepted: April 10, 2012